

PRINCIPLES OF NANO-OPTICS

LUKAS NOVOTNY

University of Rochester

BERT HECHT

University of Basel



CAMBRIDGE
UNIVERSITY PRESS

Contents

| | |
|--|----------------|
| <i>Preface</i> | <i>page xv</i> |
| 1 Introduction | 1 |
| 1.1 Nano-optics in a nutshell | 3 |
| 1.2 Historical survey | 5 |
| 1.3 Scope of the book | 7 |
| References | 11 |
| 2 Theoretical foundations | 13 |
| 2.1 Macroscopic electrodynamics | 14 |
| 2.2 Wave equations | 15 |
| 2.3 Constitutive relations | 15 |
| 2.4 Spectral representation of time-dependent fields | 17 |
| 2.5 Time-harmonic fields | 17 |
| 2.6 Complex dielectric constant | 18 |
| 2.7 Piecewise homogeneous media | 19 |
| 2.8 Boundary conditions | 19 |
| 2.8.1 Fresnel reflection and transmission coefficients | 21 |
| 2.9 Conservation of energy | 23 |
| 2.10 Dyadic Green's functions | 25 |
| 2.10.1 Mathematical basis of Green's functions | 25 |
| 2.10.2 Derivation of the Green's function for the electric field | 26 |
| 2.10.3 Time-dependent Green's functions | 30 |
| 2.11 Evanescent fields | 31 |
| 2.11.1 Energy transport by evanescent waves | 35 |
| 2.11.2 Frustrated total internal reflection | 36 |
| 2.12 Angular spectrum representation of optical fields | 38 |
| 2.12.1 Angular spectrum representation of the dipole field | 42 |

| | |
|---|------------|
| Problems | 43 |
| References | 43 |
| 3 Propagation and focusing of optical fields | 45 |
| 3.1 Field propagators | 45 |
| 3.2 Paraxial approximation of optical fields | 47 |
| 3.2.1 Gaussian laser beams | 47 |
| 3.2.2 Higher-order laser modes | 50 |
| 3.2.3 Longitudinal fields in the focal region | 50 |
| 3.3 Polarized electric and polarized magnetic fields | 53 |
| 3.4 Far-fields in the angular spectrum representation | 54 |
| 3.5 Focusing of fields | 56 |
| 3.6 Focal fields | 61 |
| 3.7 Focusing of higher-order laser modes | 66 |
| 3.8 Limit of weak focusing | 71 |
| 3.9 Focusing near planar interfaces | 73 |
| 3.10 Reflected image of a strongly focused spot | 78 |
| Problems | 86 |
| References | 87 |
| 4 Spatial resolution and position accuracy | 89 |
| 4.1 The point-spread function | 89 |
| 4.2 The resolution limit(s) | 95 |
| 4.2.1 Increasing resolution through selective excitation | 98 |
| 4.2.2 Axial resolution | 100 |
| 4.2.3 Resolution enhancement through saturation | 102 |
| 4.3 Principles of confocal microscopy | 105 |
| 4.4 Axial resolution in multiphoton microscopy | 110 |
| 4.5 Position accuracy | 111 |
| 4.5.1 Theoretical background | 112 |
| 4.5.2 Estimating the uncertainties of fit parameters | 115 |
| 4.6 Principles of near-field optical microscopy | 121 |
| 4.6.1 Information transfer from near-field to far-field | 125 |
| Problems | 131 |
| References | 132 |
| 5 Nanoscale optical microscopy | 134 |
| 5.1 Far-field illumination and detection | 134 |
| 5.1.1 Confocal microscopy | 134 |
| 5.2 Near-field illumination and far-field detection | 147 |
| 5.2.1 Aperture scanning near-field optical microscopy | 148 |
| 5.2.2 Field-enhanced scanning near-field optical microscopy | 149 |

| | | |
|----------|--|------------|
| 5.3 | Far-field illumination and near-field detection | 157 |
| 5.3.1 | Scanning tunneling optical microscopy | 157 |
| 5.3.2 | Collection mode near-field optical microscopy | 162 |
| 5.4 | Near-field illumination and near-field detection | 163 |
| 5.5 | Other configurations: energy-transfer microscopy | 165 |
| 5.6 | Conclusion | 169 |
| | Problems | 169 |
| | References | 169 |
| 6 | Near-field optical probes | 173 |
| 6.1 | Dielectric probes | 173 |
| 6.1.1 | Tapered optical fibers | 174 |
| 6.1.2 | Tetrahedral tips | 179 |
| 6.2 | Light propagation in a conical dielectric probe | 179 |
| 6.3 | Aperture probes | 182 |
| 6.3.1 | Power transmission through aperture probes | 184 |
| 6.3.2 | Field distribution near small apertures | 189 |
| 6.3.3 | Near-field distribution of aperture probes | 193 |
| 6.3.4 | Enhancement of transmission and directionality | 195 |
| 6.4 | Fabrication of aperture probes | 197 |
| 6.4.1 | Aperture formation by focused ion beam milling | 200 |
| 6.4.2 | Electrochemical opening and closing of apertures | 201 |
| 6.4.3 | Aperture punching | 202 |
| 6.4.4 | Microfabricated probes | 203 |
| 6.5 | Optical antennas: tips, scatterers, and bowties | 208 |
| 6.5.1 | Solid metal tips | 208 |
| 6.5.2 | Particle-plasmon probes | 215 |
| 6.5.3 | Bowtie antenna probes | 218 |
| 6.6 | Conclusion | 219 |
| | Problems | 220 |
| | References | 220 |
| 7 | Probe-sample distance control | 225 |
| 7.1 | Shear-force methods | 226 |
| 7.1.1 | Optical fibers as resonating beams | 227 |
| 7.1.2 | Tuning-fork sensors | 230 |
| 7.1.3 | The effective harmonic oscillator model | 232 |
| 7.1.4 | Response time | 234 |
| 7.1.5 | Equivalent electric circuit | 236 |
| 7.2 | Normal force methods | 238 |
| 7.2.1 | Tuning fork in tapping mode | 239 |
| 7.2.2 | Bent fiber probes | 240 |

| | | |
|----------|--|------------|
| 7.3 | Topographic artifacts | 240 |
| 7.3.1 | Phenomenological theory of artifacts | 243 |
| 7.3.2 | Example of near-field artifacts | 245 |
| 7.3.3 | Discussion | 246 |
| | Problems | 247 |
| | References | 248 |
| 8 | Light emission and optical interactions in nanoscale environments | 250 |
| 8.1 | The multipole expansion | 251 |
| 8.2 | The classical particle–field Hamiltonian | 255 |
| 8.2.1 | Multipole expansion of the interaction Hamiltonian | 258 |
| 8.3 | The radiating electric dipole | 260 |
| 8.3.1 | Electric dipole fields in a homogeneous space | 261 |
| 8.3.2 | Dipole radiation | 265 |
| 8.3.3 | Rate of energy dissipation in inhomogeneous environments | 266 |
| 8.3.4 | Radiation reaction | 268 |
| 8.4 | Spontaneous decay | 269 |
| 8.4.1 | QED of spontaneous decay | 270 |
| 8.4.2 | Spontaneous decay and Green’s dyadics | 273 |
| 8.4.3 | Local density of states | 276 |
| 8.5 | Classical lifetimes and decay rates | 277 |
| 8.5.1 | Homogeneous environment | 277 |
| 8.5.2 | Inhomogeneous environment | 281 |
| 8.5.3 | Frequency shifts | 282 |
| 8.5.4 | Quantum yield | 283 |
| 8.6 | Dipole–dipole interactions and energy transfer | 284 |
| 8.6.1 | Multipole expansion of the Coulombic interaction | 284 |
| 8.6.2 | Energy transfer between two particles | 285 |
| 8.7 | Delocalized excitations (strong coupling) | 294 |
| 8.7.1 | Entanglement | 299 |
| | Problems | 300 |
| | References | 302 |
| 9 | Quantum emitters | 304 |
| 9.1 | Fluorescent molecules | 304 |
| 9.1.1 | Excitation | 305 |
| 9.1.2 | Relaxation | 306 |
| 9.2 | Semiconductor quantum dots | 309 |
| 9.2.1 | Surface passivation | 310 |
| 9.2.2 | Excitation | 312 |
| 9.2.3 | Coherent control of excitons | 313 |

| | | |
|-----------|---|------------|
| 9.3 | The absorption cross-section | 315 |
| 9.4 | Single-photon emission by three-level systems | 318 |
| 9.4.1 | Steady-state analysis | 319 |
| 9.4.2 | Time-dependent analysis | 320 |
| 9.5 | Single molecules as probes for localized fields | 325 |
| 9.5.1 | Field distribution in a laser focus | 327 |
| 9.5.2 | Probing strongly localized fields | 329 |
| 9.6 | Conclusion | 332 |
| | Problems | 333 |
| | References | 333 |
| 10 | Dipole emission near planar interfaces | 335 |
| 10.1 | Allowed and forbidden light | 336 |
| 10.2 | Angular spectrum representation of the dyadic Green's function | 338 |
| 10.3 | Decomposition of the dyadic Green's function | 339 |
| 10.4 | Dyadic Green's functions for the reflected and transmitted fields | 340 |
| 10.5 | Spontaneous decay rates near planar interfaces | 343 |
| 10.6 | Far-fields | 346 |
| 10.7 | Radiation patterns | 350 |
| 10.8 | Where is the radiation going? | 353 |
| 10.9 | Magnetic dipoles | 356 |
| 10.10 | Image dipole approximation | 357 |
| 10.10.1 | Vertical dipole | 358 |
| 10.10.2 | Horizontal dipole | 359 |
| 10.10.3 | Including retardation | 359 |
| | Problems | 360 |
| | References | 361 |
| 11 | Photonic crystals and resonators | 363 |
| 11.1 | Photonic crystals | 363 |
| 11.1.1 | The photonic bandgap | 364 |
| 11.1.2 | Defects in photonic crystals | 368 |
| 11.2 | Optical microcavities | 370 |
| | Problems | 377 |
| | References | 377 |
| 12 | Surface plasmons | 378 |
| 12.1 | Optical properties of noble metals | 379 |
| 12.1.1 | Drude–Sommerfeld theory | 380 |
| 12.1.2 | Interband transitions | 381 |
| 12.2 | Surface plasmon polaritons at plane interfaces | 382 |
| 12.2.1 | Properties of surface plasmon polaritons | 386 |

| | | |
|-----------|---|-----|
| 12.2.2 | Excitation of surface plasmon polaritons | 387 |
| 12.2.3 | Surface plasmon sensors | 392 |
| 12.3 | Surface plasmons in nano-optics | 393 |
| 12.3.1 | Plasmons supported by wires and particles | 398 |
| 12.3.2 | Plasmon resonances of more complex structures | 407 |
| 12.3.3 | Surface-enhanced Raman scattering | 410 |
| 12.4 | Conclusion | 414 |
| | Problems | 414 |
| | References | 416 |
| 13 | Forces in confined fields | 419 |
| 13.1 | Maxwell's stress tensor | 420 |
| 13.2 | Radiation pressure | 423 |
| 13.3 | The dipole approximation | 424 |
| 13.3.1 | Time-averaged force | 426 |
| 13.3.2 | Monochromatic fields | 427 |
| 13.3.3 | Saturation behavior for near-resonance excitation | 429 |
| 13.3.4 | Beyond the dipole approximation | 432 |
| 13.4 | Optical tweezers | 433 |
| 13.5 | Angular momentum and torque | 436 |
| 13.6 | Forces in optical near-fields | 437 |
| 13.7 | Conclusion | 443 |
| | Problems | 443 |
| | References | 444 |
| 14 | Fluctuation-induced interactions | 446 |
| 14.1 | The fluctuation–dissipation theorem | 446 |
| 14.1.1 | The system response function | 448 |
| 14.1.2 | Johnson noise | 452 |
| 14.1.3 | Dissipation due to fluctuating external fields | 454 |
| 14.1.4 | Normal and antinormal ordering | 455 |
| 14.2 | Emission by fluctuating sources | 456 |
| 14.2.1 | Blackbody radiation | 458 |
| 14.2.2 | Coherence, spectral shifts and heat transfer | 459 |
| 14.3 | Fluctuation-induced forces | 461 |
| 14.3.1 | The Casimir–Polder potential | 463 |
| 14.3.2 | Electromagnetic friction | 467 |
| 14.4 | Conclusion | 472 |
| | Problems | 472 |
| | References | 473 |

| | |
|--|------------|
| 15 Theoretical methods in nano-optics | 475 |
| 15.1 The multiple multipole method | 476 |
| 15.2 Volume integral methods | 483 |
| 15.2.1 The volume integral equation | 484 |
| 15.2.2 The method of moments (MOM) | 490 |
| 15.2.3 The coupled dipole method (CDM) | 490 |
| 15.2.4 Equivalence of the MOM and the CDM | 492 |
| 15.3 Effective polarizability | 494 |
| 15.4 The total Green's function | 495 |
| 15.5 Conclusion and outlook | 496 |
| Problems | 497 |
| References | 498 |
| Appendix A Semianalytical derivation of the atomic polarizability | 500 |
| A.1 Steady-state polarizability for weak excitation fields | 504 |
| A.2 Near-resonance excitation in absence of damping | 506 |
| A.3 Near-resonance excitation with damping | 508 |
| Appendix B Spontaneous emission in the weak coupling regime | 510 |
| B.1 Weisskopf–Wigner theory | 510 |
| B.2 Inhomogeneous environments | 512 |
| References | 514 |
| Appendix C Fields of a dipole near a layered substrate | 515 |
| C.1 Vertical electric dipole | 515 |
| C.2 Horizontal electric dipole | 516 |
| C.3 Definition of the coefficients A_j , B_j , and C_j | 519 |
| Appendix D Far-field Green's functions | 521 |
| Index | 525 |